

Kähler Fermions on the Weitzenböck space-time .

I. B. Pestov*

*Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research
Dubna, 141980, Russia
(February 1, 2008)*

The Kähler-Dirac equation is derived on the Weitzenböck space-time, which has a quadruplet of parallel vector fields as the fundamental structure. A consistent system of equations for the Kähler fields and parallel vector fields is obtained.

Key words: teleparallelism, fermions, internal symmetry, equations.

12.20.-m, 12.20.Ds, 78.60.Mq

I. INTRODUCTION

The notion of absolute parallelism or teleparallelism was introduced by Einstein [1] when he tried to unify gravitation and electromagnetism. A transcription of the Dirac equation as a set of equations for an antisymmetric tensor field was introduced by mathematician E. Kähler [2](see also [3]). This Kähler-Dirac equation has been studied in connection with lattice fermions [4],[5] and other remarkable properties [6]-[10]. As it was suggested by Graf [6] the Kähler field might be more fundamental than the Dirac spinor. This is an appealing idea because it conforms to the Einstein methodology of associating all physical fields with geometrical objects.

Here, the Kähler-Dirac equation on the Weitzenböck space-time [11] characterized by the vanishing curvature tensor (absolute parallelism) and by the torsion tensor formed of four parallel vector fields (parallel frames) is studied. Motivation for our consideration is the known fact that orthonormal frames are necessary for description of gravitational interactions of fermions in the framework of the usual spinor formalism. To investigate the role of frames in the Kähler formalism, it is natural to consider the Kähler-Dirac equation on the Weitzenböck space-time. Our main result is the consistent system of equations (1),(2),(3),(4) for the Kähler fields and parallel vector fields with 16 degrees of freedom defined by the internal symmetry inherent in the Kähler-Dirac equation on the Weitzenböck space-time. We do not consider the physical interpretation of the equations derived because at that time it was absent for the Kähler field on a more familiar Riemann space-time, but the results of our consideration may be useful for solving this problem.

II. KÄHLER-DIRAC EQUATION ON THE WEITZENBÖCK SPACE-TIME

The Weitzenböck space-time admits a quadruplet of linearly independent parallel vector fields h_i^μ , defined by

$$\nabla_\nu h_i^\mu = \partial_\nu h_i^\mu + \Gamma_{\nu\lambda}^\mu h_i^\lambda = 0.$$

Solving these equations, we find the non-symmetric connexion,

$$\Gamma_{\mu\nu}^\lambda = h_i^\lambda \partial_\mu h_\nu^i,$$

and the torsion tensor,

$$S_{\mu\nu}^\lambda = \frac{1}{2} h_i^\lambda (\partial_\mu h_\nu^i - \partial_\nu h_\mu^i).$$

Here h_μ^i is also a quadruplet of parallel covector fields, inverse to h_i^μ . Thus, the coefficients h_i^μ or h_μ^i , are 16 functions and must satisfy the relations

$$h_i^\mu h_\nu^i = \delta_\nu^\mu, \quad h_j^\mu h_\mu^i = \delta_j^i,$$

*Electronic address: pestov@thsun1.jinr.ru

$$\eta_{ij}h_\mu^i h_\nu^j = g_{\mu\nu}, \quad \eta^{ij}h_i^\mu h_j^\nu = g^{\mu\nu},$$

where $\eta_{ij} = \eta^{ij} = \text{diag}(+, -, -, -)$.

The covariant antisymmetric tensor field $f_{\mu_1 \dots \mu_p}$ ($p = 0, 1, 2, 3, 4$) is called the p -form. If

$$F = (f, f_\mu, f_{\mu\nu}, f_{\mu\nu\lambda}, f_{\mu\nu\lambda\sigma})$$

is an inhomogeneous form, (i.e. Kähler field) the generalized curl operator D_e is defined by

$$D_e F = (0, D_\mu f, 2D_{[\mu} f_{\nu]}, 3D_{[\mu} f_{\nu\lambda]}, 4D_{[\mu} f_{\nu\lambda\sigma]}),$$

where $D_\mu = \nabla_\mu + S_\mu$, the covector S_μ is equal to the contraction of the torsion tensor $S_\mu = S_{\mu\tau}^\tau$. Square brackets denote alternation. For the operator D_i of generalized divergence we have the following definition

$$D_i F = (-D^\tau f_\tau, -D^\tau f_{\tau\mu}, -D^\tau f_{\tau\mu\nu}, -D^\tau f_{\tau\mu\nu\lambda}, 0).$$

The Kähler-Dirac equation on the Weitzenböck space-time is of the form

$$DF = \frac{mc}{\hbar} F, \quad (2.1)$$

where $D = D_i + D_e$.

Similar to the operators D_i and D_e one can introduce the operators Q_i and Q_e defined by the vector field u^μ as follows

$$Q_e F = (0, u_\mu f, 2u_{[\mu} f_{\nu]}, 3u_{[\mu} f_{\nu\lambda]}, 4u_{[\mu} f_{\nu\lambda\sigma]}),$$

$$Q_i F = (-u^\tau f_\tau, -u^\tau f_{\tau\mu}, -u^\tau f_{\tau\mu\nu}, -u^\tau f_{\tau\mu\nu\lambda}, 0).$$

If in addition to these operators, we introduce a numerical operator Λ such that

$$\Lambda F = (f, -f_\mu, f_{\mu\nu}, -f_{\mu\nu\lambda}, f_{\mu\nu\lambda\sigma}),$$

then it can be shown that the operator $Q = (Q_i - Q_e)\Lambda$ commutes with the operator D under the condition $\nabla_\mu u^\nu = 0$. Thus, the operator Q acts in the space of the solutions of equation (1) if $u^\mu = u^i h_i^\mu$, where u^i are constants.

This internal symmetry of equation (1) immediately gives the tensor

$$J^{\mu\nu} = \sum_{p=0}^4 \frac{(-1)^{p+1}}{p!} \left(\frac{1}{2} g^{\mu\nu} \bar{f}^{\tau_1 \dots \tau_p} f_{\tau_1 \dots \tau_p} + \bar{f}^{\mu\tau_1 \dots \tau_p} f_{\tau_1 \dots \tau_p}^\nu - \bar{f}^{\mu\nu\tau_1 \dots \tau_p} f_{\tau_1 \dots \tau_p} \right) + c.c.,$$

which on the solutions of equation (1) satisfies the equation

$$(\nabla_\tau + 2S_\tau)J^{\tau\mu} = 0. \quad (2.2)$$

To prove this, let

$$(F, H) = \sum_{p=0}^4 \frac{1}{p!} (\bar{f}^{\tau_1 \dots \tau_p} f_{\tau_1 \dots \tau_p}).$$

After some calculations it can be shown that

$$(DF + mF, H) - (F, DH + mH) = (\nabla_\tau + 2S_\tau)R^\tau,$$

where $R^\tau = (F, H)^\tau$ and

$$(F, H)^\tau = \sum_{p=0}^4 \frac{1}{p!} (\bar{f}_{\mu_1 \dots \mu_p} h^{\tau\mu_1 \dots \mu_p} - \bar{f}^{\tau\mu_1 \dots \mu_p} h_{\mu_1 \dots \mu_p}).$$

From this equality it follows that if F and H are solutions to equation (1), then the vector $R^\tau = (F, H)^\tau$ satisfies the equation

$$(\nabla_\tau + 2S_\tau)R^\tau = 0.$$

Substitution $H = QF$ gives

$$(F, QF)^\tau = u_\mu J^{\tau\mu},$$

which proves the statement.

III. EQUATION FOR PARALLEL VECTOR FIELDS

Since the tensor $J^{\tau\mu}$ has 16 components, one can write the consistent system of equations for 16 functions h_i^μ

$$E^{\mu\nu} = lJ^{\mu\nu} \quad (3.1)$$

provided that from the components of h_i^μ and their first and second derivatives one can construct the tensor $E^{\mu\nu}$ which satisfies the equation

$$(\nabla_\tau + 2S_\tau)E^{\tau\mu} = 0$$

identically. It is remarkable that a tensor like that really exists.

Let

$$S^{\mu\nu\lambda} = g^{\mu\tau} g^{\nu\sigma} S_{\tau\sigma}^\lambda.$$

From the commutator of covariant derivatives

$$\nabla_\tau \nabla_\sigma S^{\mu\nu\lambda} - \nabla_\sigma \nabla_\tau S^{\mu\nu\lambda} = -2S_{\tau\sigma}^\beta \nabla_\beta S^{\mu\nu\lambda}$$

it follows that

$$\nabla_\mu (\nabla_\nu S^{\mu\nu\lambda} + S_{\tau\sigma}^\mu S^{\tau\sigma\lambda}) = S^{\tau\sigma\lambda} \nabla_\mu S_{\tau\sigma}^\mu.$$

The right hand side of this equality admits following representation. The identity for general linear connexion [12]

$$R_{[\nu\tau\sigma]}^{\cdot\mu} = 2\nabla_{[\nu} S_{\tau\sigma]}^\mu - 4S_{[\nu\tau}^\beta S_{\sigma]\beta}^\mu$$

in our case ($R_{\nu\tau\sigma}^{\cdot\mu} = 0$) gives

$$\nabla_\nu S_{\tau\sigma}^\mu + \nabla_\tau S_{\sigma\nu}^\mu + \nabla_\sigma S_{\nu\tau}^\mu =$$

$$2S_{\nu\tau}^\beta S_{\sigma\beta}^\mu + 2S_{\tau\sigma}^\beta S_{\nu\beta}^\mu + 2S_{\sigma\nu}^\beta S_{\tau\beta}^\mu.$$

From this one derives by contraction

$$\nabla_\mu S_{\tau\sigma}^\mu = -\nabla_\tau S_\sigma + \nabla_\sigma S_\tau - 2S_\beta S_{\tau\sigma}^\beta.$$

Thus,

$$S^{\tau\sigma\lambda} \nabla_\mu S_{\tau\sigma}^\mu = -\nabla_\mu (2S_\nu S^{\mu\nu\lambda}) - 2S_\mu (\nabla_\nu S^{\mu\nu\lambda} + S_{\tau\sigma}^\mu S^{\tau\sigma\lambda}).$$

After substitution in to the starting equality we get

$$\nabla_\mu (\nabla_\nu S^{\mu\nu\lambda} + S_{\tau\sigma}^\mu S^{\tau\sigma\lambda} + 2S_\nu S^{\mu\nu\lambda}) =$$

$$-2S_\mu (\nabla_\nu S^{\mu\nu\lambda} + S_{\tau\sigma}^\mu S^{\tau\sigma\lambda} + 2S_\nu S^{\mu\nu\lambda}).$$

and hence the tensor

$$E^{\mu\lambda} = \nabla_\nu S^{\mu\nu\lambda} + S_{\tau\sigma}^\mu S^{\tau\sigma\lambda} + 2S_\nu S^{\mu\nu\lambda} \quad (3.2)$$

satisfies equation (2) identically.

Thus, we have proved that the system of equations (1),(2),(3),(4) is consistent.

Since the tensors $J^{\mu\nu}$ and $E^{\mu\nu}$ have dimensions cm^{-3} and respectively cm^{-2} , then the constant l in equation (3) has the dimension of length. It is natural to suppose that l is equal to the Planck length. It should be noted also that the 00-component of the tensor $J^{\mu\nu}$ is positive definite.

Thus, the connection is established between the theory of Kähler fermions and the spaces with teleparallelism, expressed by the system of equations (1),(2),(3),(4).

REFERENCES

1. Einstein, A. (1930). *Sitzungsber. preuss. Acad. Wiss., phys.-math. Kl.*, 401.
2. Kähler, E. (1962). *Rend. Mat.*, **(3-4) 21**, 425.
3. Ivanenko, D., and Landau, L. (1928). *Zeit. für Phys.*, **48**, 340.
4. Becher, P., Joos, H. (1982). *Zeit. für Phys.*, **C15**, 343.
5. Gockeler, M. (1984). *Phys. Lett.*, **B142**, 197.
6. Graf, W. (1978). *Ann. Inst. Henri Poincaré*, **A29**, 85.
7. Banks, I.M., Dothan, V., Horn, D. (1982). *Phys. Lett.*, **B117**, 413.
8. Benn, I.M., Tucker, R.W. (1982). *Phys. Lett.*, **B119**, 348.
9. Holdom, B. (1984). *Nucl. Phys.*, **B233**, 413.
10. Bullinaria, J.A. (1985). *Ann. of Phys.*, **159**, 272.
11. Weitzenböck, R. (1923). *Invariantentheorie* **XIII**, **7**. (Noordhoff, Groningen).
12. Schouten, J.A. (1954). *Ricci-Calculus*. (Berlin).